

COMPOSITIONS FOR TREATING CYSTIC FIBROSIS

Cross Reference to Related Applications

This application is a continuation-in-part of U.S. Application No. 09/587,818 filed June 06, 2000, which is a continuation-in-part of U.S. Application No. 09/249,674 filed February 12, 1999 (issued as U.S. Patent No. 6,159,447 on December 12, 2000), which is a continuation-in-part of U.S. Application No. 08/951,393 filed October 16, 1997 (issued as U.S. Patent No. 5,871,714 on February 16, 1999), both of which are incorporated herein by reference.

Field and Background of the Invention

Cystic fibrosis, a genetically inherited disease, is caused by the mutation of a gene that produces an electrolyte transfer protein. The consequence of the mutation affects a multitude of organ systems. However, the tissues that are most directly affected are those that secrete mucus or have mucus membranes. Serious adverse consequences occur with tissues that are associated with the respiratory system i.e. lungs and airway passage tissues. In response to the genetic defect, the host produces secretions to counteract the ionic imbalance. These secretions in response to the disease allow opportunistic infections to develop which causes additional fluid and mucus to infiltrate the respiratory system from the host. The additional fluid from the host in response to the infection is under the influence of the immune system. Finally, the infectious bacteria and the associated bacterial biofilm add additional mucus and fluid at the site. Furthermore, the principal bacterial pathogen associated with the opportunistic infection is *Pseudomonas aeruginosa*, which often mutates to a mucoid form that

is a prolific producer of alginate biofilm, which further exacerbates the disease condition. The result is an accumulation of copious amount of mucus, fluid and biofilm material which affects not only the respiratory system, but the entire host.

5

The current treatment of cystic fibrosis involves a dual approach to: 1) promote and facilitate the removal of mucus and secretions from the respiratory tract and; 2) control the infection that is associated with the disease. The infection, with the bacteria's production of a biofilm, obstructs the host's defenses, shields the bacteria from the killing action of antibiotics and increases the viscosity of the mucus, making it increasingly difficult for the patient to expectorate the interfering mucus.

Currently, there are limitations in achieving these objectives. Existing pharmacological agents that assist in removing mucus from the respiratory tract are not completely effective. In addition, since chronic use of antibiotics are necessary to treat the on-going infection, resistance develops, making effective antibiotic treatment increasingly difficult. Consequently, a therapeutic strategy, which is focused on both the microbial biofilm and the mucus, is an attractive treatment regimen for the disease.

Attacking and dismantling a biofilm structure is described by Budny (US 5,871,714) and Budny, et al. (US 6,159,447). They teach a composition for controlling bacterial growth and colonization by using an enzyme-anchor complex to dismantle bacterial biofilms. The result is reduced bacterial colony size and a reduction in the total number of bacteria.

Based on work done in the 1950s and 1960s and summarized by Skak et al. [Proc. Natl. Acad. Sci. 87, 9188-9192(1990)], the viscosity of the sputum from cystic fibrosis patients was reduced when the sputum was treated with the enzyme DNase. Consequently, a less viscous, easier flowing mucus is more readily expectorated.

Infections in cystic fibrosis patients of the lower respiratory tract pose a unique requirement for treatment. These infections represent a long-term, continual infection that is confounded by periodic acute infection episodes in the upper respiratory tract. Aerosolized antibiotics have been used and in certain cases, surgical interventions have been used in addition to antibiotics.

Summary of the Invention

According to one aspect of the invention, there is provided a composition for degrading biofilm structure associated with cystic fibrosis and the debris associated therewith, the composition comprising: an enzyme selected for its ability to dismantle the biofilm structure; an anchor molecule coupled to an enzyme to form an enzyme-anchor complex, the anchor molecule being selected for its ability to attach to a surface on or proximal the biofilm structure; wherein the attachment to the surface permits prolonged retention time of the enzyme-anchor complex where the biofilm structure and associated debris are present.

Bacteria, living in colonies, produce a biofilm that not only provides an environment for the bacteria to have close proximity to one another, but also protects the community from external challenges. The challenges originate from the host's immune system

and from antibiotics, which are administered to treat the infection. Consequently, dismantling the biofilm makes the colonies, as well as the individual bacteria, more susceptible to the host's defenses and more vulnerable to the action of antibiotics.

One aspect of the invention consists of one or more enzyme(s) whose specificity includes its (their) ability to degrade the polysaccharide backbone of the biofilm structure produced by *Pseudomonas aeruginosa* that occurs in the respiratory tract of cystic fibrosis patients. While these polysaccharide-degrading enzymes are hydrolytic, they are found in three major classifications: Carboxylic Ester Hydrolases (EC 3.1.1.-), Glycosidases (EC 3.2.-.-) and Lyases Acting on Polysaccharides (EC 4.2.2.-). Examples of these enzymes include, but are not limited to, acetyl esterases, polyguluronate hydrolase, polymannuronate hydrolase, alginate lyase and alginate depolymerase.

Attached to the enzyme(s), either through chemical synthetic procedures or recombinant technology, are one or more moieties that have the capability of binding, either reversibly (non-covalently) or irreversibly (covalent bonded) to a surface near the biofilm or the biofilm itself. Collectively, these moieties are called anchors. The moieties, selected to serve as anchors, can be agents or molecular species known to have an affinity for the biofilm, the surfaces near the biofilm or known binding domains. Examples of these types of anchors are listed below. The listing is not intended to be a complete list; rather, the listed examples serve to illustrate the entire class. Finally, the search for anchors can be accomplished with High Throughput Screening (HTS) of a biofilm

of either known or unknown composition with various molecular entities using a suitable assay to determine which materials have an affinity for the biofilm or surrounding surfaces.

5 These two properties: 1. an enzyme; and 2. a binding component that is connected to the enzyme, are directed at the degradation of the biofilm backbone structure.

Moieties with a Known Affinity for Biofilms

10 Concanavalin A
11 Wheat Germ Agglutinin
12 Other Lectins
13 Heparin Binding Domains
14 Elastase
15 Amylose Binding Protein
16 Ricinus communis agglutinin I (RCA I); Dilichos biflorus agglutinin (DBA); Ulex europaeus agglutinin I (UEA I).

Binding Domains from the Following Enzymes

20 Dextranucrase
21 Starch-synthesizing enzymes
22 Cellulose-synthesizing enzymes
23 Chitin-synthesizing enzymes
24 Glycogen-synthesizing enzymes
25 Pectate synthetase
26 Glycosyl transferase-binding domains (glucan-, mutan-, levan-,
27 Polygalactosyl-synthesizing enzymes
28 et al.

30 Certain agents have been described (US Patent Nos. 3,309,274,

3,624,219, 4,064,229 and 4,431,628) as indicators or disclosing agents for oral bacterial biofilms. In effect, these agents bind to the biofilm where they can be visualized either by the naked eye or with the aid of a light source with a wavelength that shows the agent's color. The purpose of these agents as described in the cited patents is to show location of the biofilm structure.

Since biofilm disclosing agents have the required property to bind to a biofilm, they make exceptionally good anchors in the enzyme-anchor complexes. Consequently, any molecular entity whose purpose is to serve as a biofilm disclosing agent can also be used as an anchor for the anchor enzyme complex to retain enzymes at or near a biofilm. The following is a list of examples of biofilm disclosing agents, which are examples of molecules that can serve as anchors. This list is only a selected list of examples and it is not intended to exclude other disclosing agents.

Examples of Biofilm Disclosing Agents

FD&C Red #3 (erythrosin)

Amaranth (Brilliant Blue)

Synthetic fluorescent dyes

D&C Green #8

D&C Red #s 19, 22 and 28

D&C Yellow #s 7 and 8

Natural fluorescent dyes

Chlorophyll dye

Carotene

FD&C Blue #1

FD&C Green #3

Hercules Green Shade 3

Merbromin
Betacyanines
Betamine
Betanin
5 Betaxanthines
Vulgaxathin
Ruthenium Red.

10 Another and second aspect of the invention consists of two or
more hydrolytic enzymes. One enzyme has the specificity to degrade
the biofilm's polysaccharide backbone structure of a biofilm; at
least one other enzyme is hydrolytic in nature, having the
capability to degrade extraneous byproducts of both the biofilm and
15 host-originated materials. These extraneous materials, host-
bacteria- and biofilm-derived byproducts include immuno-chemicals
such as proteins, polypeptides, glycoproteins, nucleic acids,
lipids, lipid complexes of sugars and proteins (lipopolysaccharides
and lipoproteins).

20 Blends and combinations of enzymes have been used for
industrial processing applications and that multiple enzymes, used
together, can remove biofilms (Johansen, C., Falholt, P. and Gram,
L. "Enzymatic Removal and Disinfections of Bacterial Biofilms."
Applied and Environmental Microbiology, Vol. 93, No. 9, September
25 1997, p. 3724-3728). As an illustrative example, alginate lyase,
pectinase, arabinase, cellulase, hemicellulase, β -glucanase and
xylanase each connected to elastase, with the elastase serving as
an anchor to the biofilms can be used to remove alginate biofilms.
Alginate biofilms ordinarily produced by *Pseudomonas aeruginosa* and
30 *Pseudomonas fluorescens*; however, the enzyme-anchor complex

described herein, will effectively remove alginate-based biofilms produced by any bacterial or fungal species, whether they act alone or in combination with one another to create the biofilm.

5 The enzymes capable of degrading proteins and polypeptides are found in classification EC 3.4.-.-. These proteinases include proteolytic enzymes, endopeptidases, peptidyl-peptide hydrolases, serine proteinases, acid proteinases and SH-proteinases. In a universal sense, all of the protein and peptide hydrolysis enzymes
10 cleave the amide linkage between adjacent amino acids in either a polypeptide or protein. Specific examples would include, but not be limited to, peptidases, carboxypeptidase, particle-bound amino
15 peptidase (EC 3.4.11.2), chymotrypsin, trypsin, cathepsin, thrombin, prothrombinase, plasmin, elastase, subtilisin, papain, ficin, asclepain, pepsin, chymosin, collagenase and those enzymes with EC 3.4.99.-, which possess proteinase activity of unknown mechanisms.

20 Many of the enzymes that hydrolyze glycoproteins (proteoglycans) and nucleic acids have not been specifically isolated and characterized, with known mechanisms of actions. Those proteinases, peptidyl-hydrolyases and nucleic acid hydrolases where the mechanism is not known are initially classified in either EC 3.-.- as hydrolases, most likely falling into EC 3.1.-, EC 3.2.-, EC 3.4.-, and EC 4.2.2.- (Lyases Acting on Polysaccharides). Some
25 currently identified enzymes that hydrolyze glycoproteins include: Peptidoglycan endopeptidase(hydrolase) (EC 3.4.99.17)
Heparin lyase(EC 4.2.2.7)
Heparatinase
30 Chitodextrinase (EC 3.2.1.14)

Chondroitin lyase (EC 4.2.2.4; EC 4.2.2.5)

Muramidase (EC 3.2.1.17)

N-Acetylmuramidase

Sialidase/Neuraminidase (EC 3.2.1.18)

5 β -N-Acetylhexosaminidase (EC 3.2.1.52)

α -N-Acetylhexosaminidase

β -N-Acetylglucosaminidase (EC 3.2.1.30)

Hyaluronoglucosidase (EC 3.2.1.35)

Hyaluronoglucuronidase (EC 3.2.1.36)

10 β -N-Acetylgalactosaminidase (EC 3.2.1.53)

β -Aspartylacetylglucosaminidase (EC 3.2.2.1)

et al.

Some identified hydrolases acting on nucleic acid material from the general class of EC 3.1.7.- to EC 3.1.31.- include, but are not limited to, the following:

exo-deoxyribonuclease I (EC 3.1.11.1)

exo-deoxyribonuclease Iii (EC 3.1.11.2)

exo-deoxyribonuclease (Lambda-induced) (EC 3.1.11.3)

20 exo-deoxyribonuclease (Phage Sp3-induced) (EC 3.1.11.4)

exo-deoxyribonuclease V (EC 3.1.11.5)

exo-deoxyribonuclease Vii (EC 3.1.11.6)

exo-ribonuclease (EC 3.1.13.-)

exo-ribonuclease (EC 3.1.14.-)

25 Venom exo-nuclease (non-specific) (EC 3.1.16.1)

Spleen exo-nuclease (non-specific) (EC 3.1.16.1)

endo-deoxyribonuclease I (EC 3.1.21.1)

endo-deoxyribonuclease Iv (EC 3.1.21.2)

endo-deoxyribonuclease (Type I specific) (EC 3.1.21.3)

30 endo-deoxyribonuclease (Type Ii specific) (EC 3.1.21.4)

(have) the capability of acting directly upon the bacteria, causing lysis of the bacterial cell wall. One or more moieties are attached to the enzymes, forming either a single unit or multiple units. The moieties are attached to the enzymes either through chemical synthetic procedures or recombinant technology to give the enzyme moiety the capability of binding either reversibly (non-covalently) or irreversibly (covalent bonded) to a surface near the biofilm or the biofilm itself. The purpose of this multi-enzyme system is directed at the degradation and removal of the biofilm with the contemporaneous bactericidal consequences for bacteria that were embedded in the biofilm's structure have become exposed due to the action of the biofilm-degrading enzyme(s).

Lysozyme has long been known to have bactericidal activity by destroying the bacterial cell wall, freeing cell wall components, which leads to cell lysis. Anchored lysozyme, along with anchored polysaccharide-degrading enzyme(s), can be used in concert to remove the polysaccharide backbone of a biofilm and then lyse the resident bacteria in a stepwise fashion. In a specific example of the removal of oral biofilms, lysozyme can be connected to amylase binding protein or the glucan-binding domain, either by coupling the lysozyme to the selected anchor or through a recombinant synthesis. The consequence of this combination is that the polysaccharide backbone is removed and the embedded bacteria are killed through cell lysis at the same time.

Lysozyme can be used in the treatment and removal of other biofilms along with the resident bacteria that may exist outside of the oral cavity. For biofilms produced by *Pseudomonas aeruginosa* and *Pseudomonas fluorescens*, lysozyme can be anchored with elastase

and used in conjunction with any one of the following biofilm-degrading enzymes: alginate lyase, pectinase, arabinase, cellulase, hemicellulase, β -glucanase and/or xylanase, each connected to elastase or some other suitable anchor.

5

This multi-enzyme, dual functionality for treating and eliminating biofilms can be used for any microorganism that produces a biofilm e.g., fungi and yeasts.

10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
66
67
68
69
70
71
72
73
74
75
76
77
78
79
80
81
82
83
84
85
86
87
88
89
90
91
92
93
94
95
96
97
98
99
100
101
102
103
104
105
106
107
108
109
110
111
112
113
114
115
116
117
118
119
120
121
122
123
124
125
126
127
128
129
130
131
132
133
134
135
136
137
138
139
140
141
142
143
144
145
146
147
148
149
150
151
152
153
154
155
156
157
158
159
160
161
162
163
164
165
166
167
168
169
170
171
172
173
174
175
176
177
178
179
180
181
182
183
184
185
186
187
188
189
190
191
192
193
194
195
196
197
198
199
200
201
202
203
204
205
206
207
208
209
210
211
212
213
214
215
216
217
218
219
220
221
222
223
224
225
226
227
228
229
230
231
232
233
234
235
236
237
238
239
240
241
242
243
244
245
246
247
248
249
250
251
252
253
254
255
256
257
258
259
260
261
262
263
264
265
266
267
268
269
270
271
272
273
274
275
276
277
278
279
280
281
282
283
284
285
286
287
288
289
290
291
292
293
294
295
296
297
298
299
300
301
302
303
304
305
306
307
308
309
310
311
312
313
314
315
316
317
318
319
320
321
322
323
324
325
326
327
328
329
330
331
332
333
334
335
336
337
338
339
340
341
342
343
344
345
346
347
348
349
350
351
352
353
354
355
356
357
358
359
360
361
362
363
364
365
366
367
368
369
370
371
372
373
374
375
376
377
378
379
380
381
382
383
384
385
386
387
388
389
390
391
392
393
394
395
396
397
398
399
400
401
402
403
404
405
406
407
408
409
410
411
412
413
414
415
416
417
418
419
420
421
422
423
424
425
426
427
428
429
430
431
432
433
434
435
436
437
438
439
440
441
442
443
444
445
446
447
448
449
450
451
452
453
454
455
456
457
458
459
460
461
462
463
464
465
466
467
468
469
470
471
472
473
474
475
476
477
478
479
480
481
482
483
484
485
486
487
488
489
490
491
492
493
494
495
496
497
498
499
500
501
502
503
504
505
506
507
508
509
510
511
512
513
514
515
516
517
518
519
520
521
522
523
524
525
526
527
528
529
530
531
532
533
534
535
536
537
538
539
540
541
542
543
544
545
546
547
548
549
550
551
552
553
554
555
556
557
558
559
560
561
562
563
564
565
566
567
568
569
570
571
572
573
574
575
576
577
578
579
580
581
582
583
584
585
586
587
588
589
590
591
592
593
594
595
596
597
598
599
600
601
602
603
604
605
606
607
608
609
610
611
612
613
614
615
616
617
618
619
620
621
622
623
624
625
626
627
628
629
630
631
632
633
634
635
636
637
638
639
640
641
642
643
644
645
646
647
648
649
650
651
652
653
654
655
656
657
658
659
660
661
662
663
664
665
666
667
668
669
670
671
672
673
674
675
676
677
678
679
680
681
682
683
684
685
686
687
688
689
690
691
692
693
694
695
696
697
698
699
700
701
702
703
704
705
706
707
708
709
710
711
712
713
714
715
716
717
718
719
720
721
722
723
724
725
726
727
728
729
730
731
732
733
734
735
736
737
738
739
740
741
742
743
744
745
746
747
748
749
750
751
752
753
754
755
756
757
758
759
760
761
762
763
764
765
766
767
768
769
770
771
772
773
774
775
776
777
778
779
780
781
782
783
784
785
786
787
788
789
790
791
792
793
794
795
796
797
798
799
800
801
802
803
804
805
806
807
808
809
810
811
812
813
814
815
816
817
818
819
820
821
822
823
824
825
826
827
828
829
830
831
832
833
834
835
836
837
838
839
840
841
842
843
844
845
846
847
848
849
850
851
852
853
854
855
856
857
858
859
860
861
862
863
864
865
866
867
868
869
870
871
872
873
874
875
876
877
878
879
880
881
882
883
884
885
886
887
888
889
890
891
892
893
894
895
896
897
898
899
900
901
902
903
904
905
906
907
908
909
910
911
912
913
914
915
916
917
918
919
920
921
922
923
924
925
926
927
928
929
930
931
932
933
934
935
936
937
938
939
940
941
942
943
944
945
946
947
948
949
950
951
952
953
954
955
956
957
958
959
960
961
962
963
964
965
966
967
968
969
970
971
972
973
974
975
976
977
978
979
980
981
982
983
984
985
986
987
988
989
990
991
992
993
994
995
996
997
998
999
1000
1001
1002
1003
1004
1005
1006
1007
1008
1009
1010
1011
1012
1013
1014
1015
1016
1017
1018
1019
1020
1021
1022
1023
1024
1025
1026
1027
1028
1029
1030
1031
1032
1033
1034
1035
1036
1037
1038
1039
1040
1041
1042
1043
1044
1045
1046
1047
1048
1049
1050
1051
1052
1053
1054
1055
1056
1057
1058
1059
1060
1061
1062
1063
1064
1065
1066
1067
1068
1069
1070
1071
1072
1073
1074
1075
1076
1077
1078
1079
1080
1081
1082
1083
1084
1085
1086
1087
1088
1089
1090
1091
1092
1093
1094
1095
1096
1097
1098
1099
1100
1101
1102
1103
1104
1105
1106
1107
1108
1109
1110
1111
1112
1113
1114
1115
1116
1117
1118
1119
1120
1121
1122
1123
1124
1125
1126
1127
1128
1129
1130
1131
1132
1133
1134
1135
1136
1137
1138
1139
1140
1141
1142
1143
1144
1145
1146
1147
1148
1149
1150
1151
1152
1153
1154
1155
1156
1157
1158
1159
1160
1161
1162
1163
1164
1165
1166
1167
1168
1169
1170
1171
1172
1173
1174
1175
1176
1177
1178
1179
1180
1181
1182
1183
1184
1185
1186
1187
1188
1189
1190
1191
1192
1193
1194
1195
1196
1197
1198
1199
1200
1201
1202
1203
1204
1205
1206
1207
1208
1209
1210
1211
1212
1213
1214
1215
1216
1217
1218
1219
1220
1221
1222
1223
1224
1225
1226
1227
1228
1229
1230
1231
1232
1233
1234
1235
1236
1237
1238
1239
1240
1241
1242
1243
1244
1245
1246
1247
1248
1249
1250
1251
1252
1253
1254
1255
1256
1257
1258
1259
1260
1261
1262
1263
1264
1265
1266
1267
1268
1269
1270
1271
1272
1273
1274
1275
1276
1277
1278
1279
1280
1281
1282
1283
1284
1285
1286
1287
1288
1289
1290
1291
1292
1293
1294
1295
1296
1297
1298
1299
1300
1301
1302
1303
1304
1305
1306
1307
1308
1309
1310
1311
1312
1313
1314
1315
1316
1317
1318
1319
1320
1321
1322
1323
1324
1325
1326
1327
1328
1329
1330
1331
1332
1333
1334
1335
1336
1337
1338
1339
1340
1341
1342
1343
1344
1345
1346
1347
1348
1349
1350
1351
1352
1353
1354
1355
1356
1357
1358
1359
1360
1361
1362
1363
1364
1365
1366
1367
1368
1369
1370
1371
1372
1373
1374
1375
1376
1377
1378
1379
1380
1381
1382
1383
1384
1385
1386
1387
1388
1389
1390
1391
1392
1393
1394
1395
1396
1397
1398
1399
1400
1401
1402
1403
1404
1405
1406
1407
1408
1409
1410
1411
1412
1413
1414
1415
1416
1417
1418
1419
1420
1421
1422
1423
1424
1425
1426
1427
1428
1429
1430
1431
1432
1433
1434
1435
1436
1437
1438
1439
1440
1441
1442
1443
1444
1445
1446
1447
1448
1449
1450
1451
1452
1453
1454
1455
1456
1457
1458
1459
1460
1461
1462
1463
1464
1465
1466
1467
1468
1469
1470
1471
1472
1473
1474
1475
1476
1477
1478
1479
1480
1481
1482
1483
1484
1485
1486
1487
1488
1489
1490
1491
1492
1493
1494
1495
1496
1497
1498
1499
1500
1501
1502
1503
1504
1505
1506
1507
1508
1509
1510
1511
1512
1513
1514
1515
1516
1517
1518
1519
1520
1521
1522
1523
1524
1525
1526
1527
1528
1529
1530
1531
1532
1533
1534
1535
1536
1537
1538
1539
1540
1541
1542
1543
1544
1545
1546
1547
1548
1549
1550
1551
1552
1553
1554
1555
1556
1557
1558
1559
1560
1561
1562
1563
1564
1565
1566
1567
1568
1569
1570
1571
1572
1573
1574
1575
1576
1577
1578
1579
1580
1581
1582
1583
1584
1585
1586
1587
1588
1589
1590
1591
1592
1593
1594
1595
1596
1597
1598
1599
1600
1601
1602
1603
1604
1605
1606
1607
1608
1609
1610
1611
1612
1613
1614
1615
1616
1617
1618
1619
1620
1621
1622
1623
1624
1625
1626
1627
1628
1629
1630
1631
1632
1633
1634
1635
1636
1637
1638
1639
1640
1641
1642
1643
1644
1645
1646
1647
1648
1649
1650
1651
1652
1653
1654
1655
1656
1657
1658
1659
1660
1661
1662
1663
1664
1665
1666
1667
1668
1669
1670
1671
1672
1673
1674
1675
1676
1677
1678
1679
1680
1681
1682
1683
1684
1685
1686
1687
1688
1689
1690
1691
1692
1693
1694
1695
1696
1697
1698
1699
1700
1701
1702
1703
1704
1705
1706
1707
1708
1709
1710
1711
1712
1713
1714
1715
1716
1717
1718
1719
1720
1721
1722
1723
1724
1725
1726
1727
1728
1729
1730
1731
1732
1733
1734
1735
1736
1737
1738
1739
1740
1741
1742
1743
1744
1745
1746
1747
1748
1749
1750
1751
1752
1753
1754
1755
1756
1757
1758
1759
1760
1761
1762
1763
1764
1765
1766
1767
1768
1769
1770
1771
1772
1773
1774
1775
1776
1777
1778
1779
1780
1781
1782
1783
1784
1785
1786
1787
1788
1789
1790
1791
1792
1793
1794
1795
1796
1797
1798
1799
1800
1801
1802
1803
1804
1805
1806
1807
1808
1809
1810
1811
1812
1813
1814
1815
1816
1817
1818
1819
1820
1821
1822
1823
1824
1825
1826
1827
1828
1829
1830
1831
1832
1833
1834
1835
1836
1837
1838
1839
1840
1841
1842
1843
1844
1845
1846
1847
1848
1849
1850
1851
1852
1853
1854
1855
1856
1857
1858
1859
1860
1861
1862
1863
1864
1865
1866
1867
1868
1869
1870
1871
1872
1873
1874
1875
1876
1877
1878
1879
1880
1881
1882
1883
1884
1885
1886
1887
1888
1889
1890
1891
1892
1893
1894
1895
1896
1897
1898
1899
1900
1901
1902
1903
1904
1905
1906
1907
1908
1909
1910
1911
1912
1913
1914
1915
1916
1917
1918
1919
1920
1921
1922
1923
1924
1925
1926
1927
1928
1929
1930
1931
1932
1933
1934
1935
1936
1937
1938
1939
1940
1941
1942
1943
1944
1945
1946
1947
1948
1949
1950
1951
1952
1953
1954
1955
1956
1957
1958
1959
1960
1961
1962
1963
1964
1965
1966
1967
1968
1969
1970
1971
1972
1973
1974
1975
1976
1977
1978
1979
1980
1981
1982
1983
1984
1985
1986
1987
1988
1989
1990
1991
1992
1993
1994
1995
1996
1997
1998
1999
2000
2001
2002
2003
2004
2005
2006
2007
2008
2009
2010
2011
2012
2013
2014
2015
2016
2017
2018
2019
2020
2021
2022
2023
2024
2025
2026
2027
2028
2029
2030
2031
2032
2033
2034
2035
2036
2037
2038
2039
2040
2041
2042
2043
2044
2045
2046
2047
2048
2049
2050
2051
2052
2053
2054
2055
2056
2057
2058
2059
2060
2061
2062
2063
2064
2065
2066
2067
2068
2069
2070
2071
2072
2073
2074
2075
2076
2077
2078
2079
2080
2081
2082
2083
2084
2085
2086
2087
2088
2089
2090
2091
2092
2093
2094
2095
2096
2097
2098
2099
2100
2101
2102
2103
2104
2105
2106
2107
2108
2109
2110
2111
2112
2113
2114
2115
2116
2117
2118
2119
2120
2121
2122
2123
2124
2125
2126
2127
2128
2129
2130
2131
2132
2133
2134
2135
2136
2137
2138
2139
2140
2141
2142
2143
2144
2145
2146
2147
2148
2149
2150
2151
2152
2153
2154
2155
2156
2157
2158
2159
2160
2161
2162
2163
2164
2165
2166
2167
2168
2169
2170
2171
2172
2173
2174
2175
2176
2177
2178
2179
2180
2181
2182
2183
2184
2185
2186
2187
2188
2189
2190
2191
2192
2193
2194
2195
2196
2197
2198
2199
2200
2201
2202
2203
2204
2205
2206
2207
2208
2209
2210
2211
2212
2213
2214
2215
2216
221

layer within a biofilm structure;

Fig. 3 is a schematic view of a magnified section of a single biofilm layer; and

Fig. 4 is a diagram of a Robbins-type flow cell to measure biofilm dynamics under various flow conditions and components that may be added to the flowing fluid.

Detailed Description of the Invention

The invention will be described using *Pseudomonas aeruginosa* by way of example. *Pseudomonas aeruginosa* was selected as an example because it is associated with the genetic-based disease of cystic fibrosis. However, the principles described in this invention apply to all biofilms, independent of the causative organism producing the biofilm structure.

Pseudomonas aeruginosa, which is a gram-negative rod, is one of many organisms found in slime residues associated with a wide variety of industrial, commercial and processing operations such as sewerage discharges, re-circulating water systems (cooling tower, air conditioning systems etc.), water condensate collections, paper pulping operations and, in general, any water bearing, handling, processing, collection etc. systems. Just as biofilms are ubiquitous in water handling systems, it is not surprising that *Pseudomonas aeruginosa* is also found in association with these biofilms. In many cases, *Pseudomonas aeruginosa* is the major microbial component.

In addition to its importance in industrial processes, *Pseudomonas aeruginosa* and its associated biofilm structure have far-reaching medical implications, being the basis of many

pathological conditions. *Pseudomonas aeruginosa* is an opportunistic bacterium that is associated with a wide variety of infections. It has the ability to grow at temperatures higher than many other bacteria and it is readily transferred from an environmental setting to become host-dependent. Translocation, both within a specific medium and to other media, is facilitated with its single polar flagella.

Pseudomonas aeruginosa has nutritional versatility in being able to use a wide variety of substrates, fast growth rate, motility, temperature resiliency and short incubation periods all of which contribute to its predominance in natural microflora communities as well as being the cause of nosocomial (hospital acquired) infections.

Infections caused by *Pseudomonas aeruginosa* begin usually with bacterial attachment to and colonization of mucosal and cutaneous tissues. The infection can proceed via extension to surrounding structures or infection may lead to bloodstream invasion, dissemination and sepsis syndrome. Virtually all infections that are caused by *Pseudomonas aeruginosa* as the principal cause, or as a confounding factor to other bacterial-based infections, are treatable. The one notable exception is cystic fibrosis where the infection is chronic.

Respiratory Infections. Alginate producing strains of *Pseudomonas aeruginosa* infect the lower respiratory tract of patients with cystic fibrosis leading to acute and the chronic progression of the pathological condition. In addition to cystic fibrosis, primary pneumonia often presents bilateral

bronchopneumonia with nodular infiltrates. Accompanying such infections are pleural effusions along with pathological progression leading to alveolar necrosis, focal hemorrhages and micro-abscesses.

5

Mucoid strains *Pseudomonas aeruginosa* typically infect the lower respiratory tract of individuals with cystic fibrosis. Airway obstruction typically begins with bronchial infection and mucus production followed by colonization of *Pseudomonas aeruginosa* in the lower respiratory tract. The colonization of *Pseudomonas aeruginosa* accelerates disease pathology resulting in increased mucus production, airway obstruction, bronchiectasis and fibrosis in the lungs. These conditions eventually lead to pulmonary disease leading to hypertension and hypoxemia.

It is reasonable to expect a reduction in viscosity of the mucus if the biofilm produced by the bacteria were dismantled. Since the biofilm, can and likely does to a large measure, adhere to lung tissue surfaces, an enzyme that dismantles the biofilm, being equipped with an anchor would be effective in treating cystic fibrosis patients. The treatment with the enzyme-anchor complex would accomplish several key objectives for an effective treatment regime: 1) viscosity reduction of the sputum; 2) bacterial colony size reduction; and 3) bacteria exposition by reducing the biofilm so that antibiotics and the host's own immune system would be more effective.

Equipping DNase with an anchor and combining it with and enzyme-anchor complex having an enzyme that degrades the biofilm would allow greater retention of these two, though functionally

different enzymes, for the treatment of cystic fibrosis.

While cystic fibrosis is a chronic infection of *Pseudomonas aeruginosa*, other, acute, respiratory infections occur as a result of bacteria other than *Pseudomonas aeruginosa*. For example, *Streptococcus pyogenes* is the primary cause of bacterial pharyngitis which, if uncontrolled, can lead to rheumatic fever. Nelson, et al. [Proc. Acad. Sci. 98, 4107-4112(2001)] report a lysis process to control the bacterial infection using double-stranded DNA bacteriophages. The enzymes associated with the bacteriophage-mediated lysis serve as examples of implementing the present invention.

Alginate Biofilms of *Pseudomonas aeruginosa*: At the root of *Pseudomonas aeruginosa* initial colonization, as well as its proliferative growth rate, is the production of a mucoid exopolysaccharide layer comprised of alginate. This exopolysaccharide layer, along with lipopolysaccharide, protects the organism from direct antibody and complement mediated bactericidal mechanisms and from opsonophagocytosis. This protective biofilm allows *Pseudomonas aeruginosa* to expand, grow and to exist in harsh environments that may exist outside the alginate biofilm. It is not surprising that the alginate biofilm is considered as an important virulence factor.

The alginate biofilm or "slime matrix" consists of a secreted polysaccharide that serves as the backbone structure of the biofilm. Alginate is a polysaccharide copolymer of β -D-mannuronic acid and α -L-guluronic acid linked together by 1-4 linkages. The immediate precursor to the biosynthetic polymerization is guanosine

5'-diphosphate-mannuronic acid, which is converted to mannuronan. Post-polymerization of the mannuronan by acetylation at O-2 and O-3 and epimerization, principally at C-5, of some of the monomeric units to produce gulonate, results in varying degrees of acetylation and gulonate residues. Both the degree of acetylation and the percentage of mannuronic residues that have been converted to gulonate residues greatly affect the properties of the biofilm. For example, polymers rich in gulonate residues and in the presence of calcium, tend to be more rigid and stiff than polymers with low levels of gulonate monomeric units.

Construction of Anchor-Enzyme Complexes.

The Anchor Enzyme Complex can be constructed using chemical synthetic techniques. Additionally, the Anchor-Enzyme Complex, if the anchor is a polypeptide or protein, such as protein binding domains, lectins, selectins, heparin binding domains etc., can be constructed using recombinant genetic engineering techniques.

Types of Anchors.

The binding domain from elastase

Domains that bind to carbohydrates and polysaccharide

Lectins

Mannose Binding Lectin

Selectins

The binding domains from Heparin

The binding domains of Fibronectin

CD44 Protein

Additional anchors listed in US Patent No. 5,871,714.

Type of enzymes

1. Generally, enzymes in the class EC 4.2.2._, which are polysaccharide lyases, which degrade the polysaccharide backbone

structure of biofilms:

Glycoside Hydrolases, Galactoaminidases, Galactosidases,
Glucosaminidases, Glucosidases, Mannosidases (EC 3.1.2.-)

Neuraminidase (EC 3.1.2.18)

5 Dextranase, Mutanase, Mucinase, Amylase, Fructanase,
Galactosidase, Muramidase, Levanase, Neuraminidase (EC 3.2.-)

α -Glucosidases (EC 3.2.1.20)

β -Glucosidase (EC 3.2.1.21)

α -Glucosidase (EC 3.2.1.22)

10 β -D-Mannosidase (EC 3.2.1.25)

Acetylglucosaminidase (EC 3.2.1.30)

Hyaluronoglucosaminidase (EC 3.2.1.35)

α -L-Fucosidase (EC 3.2.1.51)

Hyaluronate Lyase (EC 4.2.2.1)

15 Pectate Lyase (EC 4.2.2.2)

Alginate Lyase [Poly(β -D-Mannuronate) Lyase] (EC 4.2.2.3)

Chondroitin ABC Lyase (EC 4.2.2.4)

Chondroitin AC Lyase (EC 4.2.2.5)

Oligogalacturonide Lyase (EC 4.2.2.6)

20 Heparin Lyase (EC 4.2.2.7)

Heparan Lyase [Heparitin-Sulfate Lyase] (EC 4.2.2.8)

Exopolygalacturonate Lyase (EC 4.2.2.9)

Pectin Lyase (EC 4.2.2.10)

Poly (α -L-Guluronate) Lyase (EC 4.2.2.11)

25 Xanthan Lyase (EC 4.2.2.12)

Exo-(1,4)- α -D-Glucan Lyase (EC 4.2.2.13)

Non-specific polysaccharide depolymerases derived from
bacteriophages et al.

2. Enzymes for removing debris either embedded within the biofilm structure or extraneous byproducts of both the biofilm and host-originated materials. This later debris may originate from the host and would include immune response products. These include many EC sub-classes with the general class of hydrolytic and digestive enzymes. In descriptive terms, they include enzymes that facilitate the breaking of chemical bonds and include the following:

Esterases - cleavage of ester bonds;

Glycolytic - cleavage of bonds found in oligo- and polysaccharides

Peptidases - cleavage of peptide bonds where the substrate is a protein or polypeptide;

Nucleic acid materials (RNA and DNA);

Carbon-nitrogen cleavage - where the substrate is not a protein or polypeptide;

Acid anhydride cleaving enzymes;

Carbon-carbon bond cleavage;

Halide bond cleavage;

Phosphorus-nitrogen bond cleavage;

Sulfur-nitrogen bond cleavage; and

Carbon-phosphorus bond cleavage.

Typical examples include the following enzymes:

Endopeptidases; Peptide Hydrolases (EC 3.4.-)

Aminopeptidases (EC 3.4.11)

Nucleic Acid Hydrolases (EC 3.1.-.-)

Propyl Aminopeptidases (EC 3.4.11.5)

Glycylpropyl Dipeptidases; Dipeptidyl Peptidase (EC 3.4.14)

Serine Endopeptidases (EC 3.4.21)

Chymotrypsin (EC 3.4.21.1)

Trypsin (EC 3.4.21.4)

Amidohydrolases (EC 3.5.-)

N-Acetylglucosamine-6-phosphate Deacetylase (EC 3.5.1.25)

Oxo-Acid Lyases (EC 4.1.3.-)

N-Acetylmuraminate Lyases (EC 4.1.3.3)

5 Carbohydrate Epimerases (EC 5.1.3-)

Glucosamine-6-phosphate Isomerases (EC 5.3.1.10)

Example

10 Since *Pseudomonas aeruginosa* is a ubiquitous bacterial strain,
found not only in the environment and in industrial settings where
fouling occurs, but also in many disease conditions, it will serve
as an example to illustrate the principles of the invention.
Further, while there are many disease conditions for which
15 *Pseudomonas aeruginosa* is the cause, ocular infections will
exemplify the implementation of the invention. The choice of
Pseudomonas aeruginosa as the biofilm-producing bacteria and
pathogen and ocular infection as a consequence of the biofilm is
not meant to preclude or limit the scope of this invention. The
principles outlined in this example readily apply to all biofilms,
20 whether produced by bacteria or other organisms, all biofilms that
are generated by organisms and the embodiments, taken and
implemented either individually or collectively.

25 *Pseudomonas aeruginosa* is an opportunistic bacterial species,
which once colonized at a site such as ocular tissue, produces a
biofilm with a polysaccharide-based alginate polymer. This
exopolysaccharide or glycocalyx matrix is the confine in which the
bacterial species can grow and proliferate. This biofilm matrix can
also serve as a medium for other, pathogenic bacteria, fungi and
30 viruses. It is of therapeutic benefit, therefore, to remove the

biofilm structure and eliminate all pathogens at the site, not only *Pseudomonas aeruginosa*.

Alginate lyase, the expression product from the algL gene, can be obtained from various bacterial sources e.g. *Azotobacter vinelandii*, *Pseudomonas syringe*, *Pseudomonas aeruginosa* etc., producing an enzyme AlgL, which degrades alginate. Other genes, e.g. alxM, also provide a wide variety of alginate lyase and polysaccharide depolymerase enzymes with degrade alginate by various mechanisms.

Endogenous lectins, heparin binding domains and various receptors from animals and plants have receptors that bind to alginate. These receptors, when located on host cell surfaces, allow the evolving alginate biofilm to be retained by the infected tissue. Elastase (Leukocyte Elastase, EC 3.4.21.37 and Pancreatic Elastase, EC 3.4.21.36), which is a digestive enzyme, also has a domain that binds to alginate. Such binding capability, along with the degradative ability of the catalytic site in elastase, has been implicated in tissue degradation associated with alginate biofilm infections such as cystic fibrosis. In addition, other serine proteases also have alginate binding domains.

In one aspect of the invention, a fusion protein is created, using standard genetic engineering techniques. One of the traits or elements of the fusion protein is the ability to degrade alginate and a second property being a binding capability of the newly-created fusion protein, derived from, for example, the binding domain of elastase. The bi-functional protein fulfills the criteria set out in the invention in that the binding domain derived from

elastase serves as the anchor and the alginate lyase portion of the fusion protein serves as the degradative enzyme for the biofilm.

It is within the scope of this invention that the principles outlined here also apply to all biofilms in all circumstances in which they occur.

Construction of the Enzyme Anchor Complex

Using molecular biology and biotechnology techniques, gene fusions are created to produce unique proteins from recombinant DNA segments. A DNA sequence, which specifically codes for an enzyme is fused to a DNA segment that specifically codes for a protein-binding domain. The resulting fused DNA segment will produce a unique protein that possesses both enzymatic or catalytic activity and binding activity.

The DNA sequence that codes for alginate lyase obtained from *Pseudomonas aeruginosa*, or another acceptable strain, was isolated and amplified using polymerase chain reaction. The sequence was subcloned into an expression vector. Next the DNA that codes for leukocyte elastase was isolated from a mouse complimentary DNA (cDNA) library. The mouse leukocyte elastase sequence was amplified by using polymerase chain reaction.

Both DNA sequences for alginate lyase and mouse leukocyte elastase were subcloned into a single open reading frame within a suitable expression vector. Thus, yielding a DNA sequence that codes for a single protein that contains both the amino acid sequence for alginate lyase as well as the sequence for leukocyte elastase. This hybrid or chimeric protein has the catalytic ability

to degrade alginate as well as the binding ability of elastase.

Assay Procedure for Synthesized Anchor Enzyme Complexes

Preparation of Bacterial Biofilms. There are many procedures to prepare bacterial biofilms. Herein are two of those procedures.

Analysis of a Dynamic System. The appropriate bacterial strain, or mixed strains if more than one strain is used, is incubated in tryptic soy broth for 18 to 24 hours at 37°C. After the incubation period, the cells are washed three times with isotonic saline and re-suspended in isotonic saline to a density of 10⁶ CFU/ml. The re-suspended cells are incubated a second time with Teflon squares (1 x 1 cm) with a thickness of 0.3 cm for six to seven days at 37° C. The recovered cells in the saline incubation medium are planktonic bacteria, while those associated with the Teflon squares and the biofilm are sessile cells.

The biofilm-associated sessile cells are then treated with appropriate anchor-enzyme complexes that degrade the generated biofilm at various concentrations with or without bactericidal agents in either a completely closed system or an open system (flow-through chamber or cell). The bactericidal agent can be either an anchor enzyme system that generates active oxygen or a non-enzymatic, chemical that is a recognized antimicrobial agent, biocide or antibiotic.

Analysis of a Static System. The Teflon squares with the associated biofilm are transferred to isotonic saline medium containing a given concentration of anchor-enzyme complex that degrades the biofilm. At intervals of 3, 6, 12, 24 and 48 hours, the individual Teflon squares are washed three times with isotonic saline and finally added to fresh isotonic saline which is

vigorously shaken or sonicated for tow minutes. The suspended mixture is diluted and counted for cell density and expressed as number of CFU/ml.

5 The same counting procedure can be used for the incubation medium.

Bactericidal agents are also incorporated into the experimental design, which also uses the same cell counting procedure.

10 Estimating Biofilm Size. At the end of any of the incubation steps, the biofilm can be recovered, dehydrated and weighed to obtain total biomass of the biofilm. Alternatively, the amount of alginate backbone can be determined where the biofilm contains Pseudomonas sp.

15 Extraction of Polysaccharide Backbone. After the second incubation and disruption of the biofilm, the bacterial cells are removed from the dispersion. With an increasing concentration of an ethanol/soling gradient, the alginate is precipitated, collected and washed three times with 95% ethanol. The precipitate is desiccated after which the quantity can be determined gravimetrically or by any number of chemical, enzymatic or combination of chemical and enzymatic methods. The most widely used method is the chemical method of which there are three types:
20 uronic acid assay, orcinol-FeCl₃ and decarboxylation and CO₂ measurement.
25

30 Analysis in a Dynamic System (Complete or Partial). The most widely used dynamic flow system that can be regulated from a completely closed to a completely open system is the Robbins Device or the Modified Robbins Device. The Modified Robbins Device allows

the assessment of biofilms in which the fluid flow and growth rates of the biofilm can be regulated independently and simultaneously. A Robbins-type flow cell can be a completely closed system that possesses flow dynamics for assessing efficacy of anchor-enzyme complexes.

5

0907597-060601